

Supplementary Materials: Theoretical Simulation of the Temporal Behavior of Bragg Diffraction Derived from Lattice Deformation^{*}

Cong Guo(过聪), Shuai-Shuai Sun(孙帅帅), Lin-Lin Wei(尉琳琳),

Huan-Fang Tian(田焕芳), Huai-Xin Yang(杨槐馨), Shu Gao(高舒),

Yuan Tan(谭媛), Jian-Qi Li(李建奇)**

^{*}Supported by the National Key Research and Development Program of China under Grant Nos. 2016YFA0300303, 2017YFA0504703, and 2017YFA0302904, the National Basic Research Program of China under Grant No. 2015CB921304, the National Natural Science Foundation of China under Grant Nos. 11604372, 11474323, and 11774391, 11774403 and 61575085, the Strategic Priority Research Program (B) of the Chinese Academy of Sciences under Grant No. XDB25000000, and the Scientific Instrument Developing Project of the Chinese Academy of Sciences under Grant No. ZDKYYQ20170002.

^{**}Corresponding author. Email: ljq@iphy.ac.cn




FIG. S1 Resonance dynamics in graphite diffraction. Temporal change in the intensity integral of the graphite $(\bar{1}1\bar{1})$ Bragg spot. [1]




FIG. S2 Kinetics of the (220)-1 spot (circles) along with a fit that includes a biexponential rise component and two damped sinusoidal functions (solid line). [2]

References

- [1] Park H. S., Baskin J. S., Barwick B., Kwon O. H., and Zewail A. H., 2009 *Ultramicroscopy* **110** 7
- [2] Harb M., Peng W., Sciaini G., Hebeisen C. T., Ernstorfer R., Eriksson M. A., Lagally M. G., Kruglik S. G., and Miller R. D., 2009 *Phys. Rev. B* **79** 094301